

Dynamics and Endogeny for recursive processes on trees

JON WARREN

Department of Statistics, University of Warwick, Coventry CV4 7AL, UK.

Abstract

We consider a stochastic process $(\xi_u; u \in \Gamma_\infty)$ where Γ_∞ is the set of vertices of an infinite binary tree which satisfies the recursion relation

$$\xi_u = \phi(\xi_{u0}, \xi_{u1}, \epsilon_u) \text{ for each } u \in \Gamma_\infty.$$

Here $u0$ and $u1$ denote the two immediate daughters of the vertex u . The random variables $(\epsilon_u; u \in \Gamma_\infty)$, which are to be thought of as innovations, are supposed independent and identically distributed. A recent paper of Aldous and Bandyopadhyay has drawn attention to the issue of endogeny: that is whether the process $(\xi_u; u \in \Gamma_\infty)$ is measurable with respect to the innovations process. Here we restrict attention to the case where each ξ_u takes values in a finite set S and show how this question is related to the existence of certain dynamics. Using this we develop a necessary and sufficient condition for endogeny in terms of the coupling rate for a Markov chain on S^2 for which the diagonal is absorbing.

1 Introduction

We suppose that we are given probability spaces $(S, \mathcal{S}, \mu)$ and $(E, \mathcal{E}, \nu)$ and a mapping

$$(1.1) \quad \phi : S \times S \times E \rightarrow S$$

under which the image of the product measure $\mu \otimes \mu \otimes \nu$ is μ . In general $\phi : S^n \times E \rightarrow S$ may be considered, even with n being infinity, but here we take $n = 2$ for simplicity and we further assume that ϕ is symmetric in its first two arguments. Typically in applications ν and ϕ are given and the measure μ is to be determined. μ is then said to be a solution to a recursive distributional equation. Existence and uniqueness of μ is well studied in a number of important cases; the survey paper by Aldous and Bandyopadhyay, [1], describes many examples. Here our concerns are somewhat different. Associated with this setup is a recursive tree process constructed as follows. Let $\Gamma_\infty = \cup_{k \geq 0} \{0, 1\}^k$ be the set of vertices of the infinite rooted binary tree; in particular $\emptyset \in \Gamma_\infty$ denotes the root. The set of vertices on level n of the tree, $\{0, 1\}^n$, is denoted by G_n , and $\Gamma_n = \cup_{k=0}^n \{0, 1\}^k$ denotes the set of vertices belonging to levels up to and including n . A vertex $u = (u_1, u_2, \dots, u_n)$ has two daughters: vertices $u0 = (u_1, \dots, u_n, 0)$ and $u1 = (u_1, \dots, u_n, 1)$. Let $(\xi_u; u \in \Gamma_\infty)$ and $(\epsilon_u; u \in \Gamma_\infty)$ be the co-ordinate maps on $S^{\Gamma_\infty} \times E^{\Gamma_\infty}$. Define

$$(1.2) \quad \Omega = \{\omega \in S^{\Gamma_\infty} \times E^{\Gamma_\infty} : \xi_u(\omega) = \phi(\xi_{u0}(\omega), \xi_{u1}(\omega), \epsilon_u(\omega)) \text{ for each } u \in \Gamma_\infty\},$$

and let $\mathcal{F}$ denote the restriction of the product σ -algebra to Ω . There is a unique probability measure m on $(\Omega, \mathcal{F})$ under which for each n , the joint law of $(\xi_u; u \in G_n)$

and $(\epsilon_u; u \in \Gamma_{n-1})$ is $\otimes^{G_n} \mu \otimes^{\Gamma_{n-1}} \nu$. We think of independent random variables $(\epsilon_u; u \in \Gamma_\infty)$, sometimes referred to as the innovations, as the input that drives the system, while $(\xi_u; u \in \Gamma_\infty)$ is thought of as the response. Aldous and Bandyopadhyay draw attention to investigating when the recursive tree process is endogenous, that is to say every random variable on $(\Omega, \mathcal{F})$ is m -almost surely equal to a function of the innovations process $(\epsilon_u; u \in \Gamma_\infty)$ alone. Loosely speaking this means there is no additional randomness in the system “located” at the boundary of the infinite tree. However care must be taken in interpreting this: the tail σ -algebra

$$(1.3) \quad \bigcap_n \sigma(\xi_u, \epsilon_u; u \in \Gamma_\infty \setminus \Gamma_n)$$

is typically empty even if endogeny does not hold! A strong parallel may be drawn with certain stochastic differential equations which admit weak but not strong solutions, see in particular Tsirelson’s example discussed in Section V.18 of [9]. In non-endogenous cases it is natural to try to give some explicit description of the additional randomness, however this does not seem possible in any generality. One case that is amenable to such an analysis is the linear “smoothing transformation” which has been extensively studied. Various representations theorems obtained by Durrett and Liggett, [5], Liu [6], Caliebe and Rösler [4], and others reveal a structure in which the innovations are augmented with Gaussian or Poisson noise living on the boundary of the tree. Another case whose structure can similarly be understood is studied by Biggins [3].

Let $u_0 = \emptyset, u_1, u_2, \dots, u_n, \dots$ be an infinite sequence of vertices, u_{n+1} being a daughter of u_n for each n . For $n \leq 0$ let $\xi_n = \xi_{u_{-n}}$. The law of the sequence $(\xi_n; n \leq 0)$, which by the symmetry of ϕ does not depend on the choice of sequence of vertices, is easily seen to be that of a stationary Markov chain (indexed by negative time) with transition kernel P defined on $S \times \mathcal{S}$ given by

$$(1.4) \quad P(x_0, A) = \int_S \int_E \mathbf{1}(\phi(x_0, x_1, z) \in A) \nu(dz) \mu(dx_1).$$

In this paper we give a criterion for endogeny in terms of the corresponding “two point motion”, that is to say a Markov chain on S^2 with transition kernel $P^{(2)}$ on $S^2 \times \mathcal{S}^2$ given by

$$(1.5) \quad P^{(2)}((x_0, x'_0), A \times A') = \int_S \int_E \mathbf{1}(\phi(x_0, x_1, z) \in A, \phi(x'_0, x_1, z) \in A') \nu(dz) \mu(dx_1).$$

Let $S^\nearrow$ be the diagonal of S^2 which is an absorbing set for $P^{(2)}$. Let $P^{(-)}$ be the restriction of $P^{(2)}$ to $S^2 \setminus S^\nearrow$.

We now assume that S is finite.

In this case we can identify $P^{(-)}$ with a non-negative square matrix to which we can apply Perron-Frobenius theory. Let ρ be the largest eigenvalue of $P^{(-)}$.

Theorem 1. *If $2\rho < 1$ then the tree process is endogenous, whereas if $2\rho > 1$ then the process is not endogenous. In the critical case $2\rho = 1$, if the additional conditions (4.1) and (4.2) hold, then the tree process is endogenous.*

Aldous and Bandyopadhyay have given a different necessary and sufficient condition for endogeny, the above result being related to their condition by linearization. We now elucidate this connection. Associated with $\phi : S \times S \times E \rightarrow S$ is the “two-point map” $\phi^{(2)} : S^2 \times S^2 \times E \rightarrow S^2$ given by

$$(1.6) \quad \phi^{(2)}((x_0, x'_0), (x_1, x'_1), z) = (\phi(x_0, x_1, z), \phi(x'_0, x'_1, z)).$$

Now we can define $T^{(2)} : \mathcal{M}(S^2) \times \mathcal{M}(S^2) \rightarrow \mathcal{M}(S^2)$ by setting $T^{(2)}(\lambda_0, \lambda_1)$ equal to the image of the measure $\lambda_0 \otimes \lambda_1 \otimes \nu$ under the map $\phi^{(2)}$. Recall that μ denotes our given measure on S that is invariant for the recursive distributional equation corresponding to ϕ and ν . The measure $\mu^\nearrow$ on S^2 , carried by the diagonal and having marginals μ , is a fixed point of $T^{(2)}$ in that $T^{(2)}(\mu^\nearrow, \mu^\nearrow) = \mu^\nearrow$. According to Theorem 11 of [1], under some minor technical condition, $\mu^\nearrow$ being the only probability measure on S^2 that is a fixed point of $T^{(2)}$, and whose marginals are both equal to μ , is a necessary and sufficient condition for endogeny. Bandyopadhyay, [2], has used this criterion to show the endogeny of certain important examples that arise from applications. The map $T^{(2)}$ is bilinear and symmetric so that we have for real δ and $\lambda \in \mathcal{M}(S^2)$

$$(1.7) \quad T^{(2)}(\mu^\nearrow + \delta\lambda, \mu^\nearrow + \delta\lambda) = \mu^\nearrow + 2\delta T^{(2)}(\lambda, \mu^\nearrow) + \delta^2 T^{(2)}(\lambda, \lambda),$$

which shows that $2T^{(2)}(\cdot, \mu^\nearrow) : \mathcal{M}(S^2) \rightarrow \mathcal{M}(S^2)$ is the derivative of $\lambda \mapsto T^{(2)}(\lambda, \lambda)$ at the fixed point $\mu^\nearrow$. It is straightforward to verify that

$$(1.8) \quad \lambda P^{(2)} = T^{(2)}(\lambda, \mu^\nearrow).$$

Thus the criterion given in Theorem 1 can be viewed in terms of the stability of the fixed point $\mu^\nearrow$ of $T^{(2)}$, and it is quite natural that the existence of other fixed points of $T^{(2)}$ should be related to this.

Let $\mathcal{H} = L^2(\Omega, \mathcal{F}, m)$, and $\mathcal{K} \subseteq \mathcal{H}$ be the subspace of (equivalence classes of) random variables measurable with respect to the innovations process $(\epsilon_u; u \in \Gamma_\infty)$. Endogeny means $\mathcal{H} = \mathcal{K}$. We also have subspaces $\mathcal{K}_n$ and $\mathcal{H}_n$ containing random variables measurable with respect to $(\epsilon_u; u \in \Gamma_{n-1})$ and with respect to $(\epsilon_u; u \in \Gamma_{n-1})$ together with $(\xi_u; u \in G_n)$ respectively.

In the following three sections we treat the three cases: subcritical $2\rho < 1$; supercritical $2\rho > 1$; and critical $2\rho = 1$ separately. The subcritical case is treated by using an operator version of Markov's inequality to bound from above the quantity $\|f\|^2 - \|P_{\mathcal{K}_n} f\|^2$ where $P_{\mathcal{K}_n} f$ is the orthogonal projection of f onto $\mathcal{K}_n$. The supercritical case is treated by using the left eigenvector of $P^{(-)}$ corresponding to ρ to construct a consistent family of quadratic forms on $\mathcal{H}$, and hence an operator Q_∞ . This operator is the generator of dynamics that fix the innovations while perturbing the additional randomness at the boundary. The existence of such dynamics precludes endogeny.

In the final section of the paper we show that the dynamics associated with Q_∞ naturally arise through a passage to the limit. Consider for each $n \geq 1$ the finite configuration space

$$(1.9) \quad \Omega_n = \{\omega \in S^{\Gamma_n} \times E^{\Gamma_{n-1}} : \xi_u(\omega) = \phi(\xi_{u0}(\omega), \xi_{u1}(\omega), \epsilon_u(\omega)) \text{ for each } u \in \Gamma_{n-1}\},$$

which we equip with the measure m_n under which the joint law of $(\xi_u; u \in G_n)$ and $(\epsilon_u; u \in \Gamma_{n-1})$ is $\otimes^{G_n} \mu \otimes^{\Gamma_{n-1}} \nu$. Consider the dynamics on Ω_n with invariant

measure m_n under which the co-ordinates ξ_u for $u \in G_n$ are independently refreshed at rate 1, whilst ξ_v for $v \in \Gamma_{n-1}$ are determined by application of the map ϕ using the innovations $(\epsilon_u; u \in \Gamma_{n-1})$ which are held fixed for all time. Let $-A_n$ denote the corresponding generator acting on $L^2(\Omega_n, m_n)$. We show that at the level of generators, these dynamics, when slowed down by a factor of $(2\rho)^n$, converge to those associated with Q_∞ . Let $P_{\mathcal{H}_n} : \mathcal{H} \rightarrow \mathcal{H}_n$ be the orthogonal projection onto $\mathcal{H}_n$ which we identify with $L^2(\Omega_n, m_n)$.

Theorem 2. *Suppose that $P^{(-)}$ is primitive, and that $2\rho > 1$, then*

$$(2\rho)^{-n} A_n P_{\mathcal{H}_n} \rightarrow -Q_\infty,$$

in the strong resolvent sense as n tends to infinity.

There is strong similarity between the methods used in this paper and those used to study multitype branching processes. To make this connection fix $f \in \mathcal{H}_0$ that satisfies $\|f\|^2 = 1$ and consider the probability distribution distributions $\mu_f^{(n)}$ on the non-negative integers defined by

$$(1.10) \quad \mu_f^{(n)}(k) = \|P_{E_k} f\|^2 \text{ for } k \geq 0,$$

where P_{E_k} is the projection operator associated with the eigenspace E_k of the operator A_n corresponding to the eigenvalue k . See below at (2.3) for further information on this. We use arguments that are based on treating this distribution as if it were that of the number of particles alive in generation n of a branching process. In particular Theorem 1 corresponds exactly to the fact the certainty or otherwise of eventual extinction for a branching process depends upon whether the Malthusian parameter of the process is greater than one. The arguments used are also closely related to the spectral methods employed by Tsirelson, [10], to study continuous products of probability spaces. In fact Tsirelson and Vershik introduced certain recursive tree processes in [11] as examples in the theory of continuous products.

2 The subcritical case: endogeny

Recall the probability measure m on $(\Omega, \mathcal{F})$ is characterized by the fact that the joint law of $(\xi_u; u \in G_n)$ and $(\epsilon_u; u \in \Gamma_{n-1})$ is $\otimes^{G_n} \mu \otimes^{\Gamma_{n-1}} \nu$. Thus there are isomorphisms between Hilbert spaces:

$$(2.1) \quad \begin{aligned} \mathcal{H}_n &\cong L^2(S^{G_n} \times E^{\Gamma_{n-1}}, \mathcal{S}^{G_n} \times \mathcal{E}^{\Gamma_{n-1}}, \otimes^{G_n} \mu \otimes^{\Gamma_{n-1}} \nu) \\ &\cong \bigotimes^{G_n} L^2(S, \mathcal{S}, \mu) \bigotimes^{\Gamma_{n-1}} L^2(E, \mathcal{E}, \nu). \end{aligned}$$

$L^2(S, \mathcal{S}, \mu)$ is the direct sum of the one-dimensional subspace of constants together with its orthogonal complement to be denoted by $L_0^2(S, \mathcal{S}, \mu)$. Decomposing each copy of $L^2(S, \mathcal{S}, \mu)$ appearing on the righthandside of (2.1) in this manner we obtain

$$(2.2) \quad \mathcal{H}_n = \bigoplus_{S \subseteq G_n} \mathcal{H}_S,$$

where for each subset $S \subseteq G_n$ of vertices on level n of the tree the corresponding subspace $\mathcal{H}_S$ of the Hilbert space $\mathcal{H}_n$ is generated by vectors of the form $\otimes_{u \in G_n} f_u \otimes_{v \in \Gamma_{n-1}} g_v$ with $f_u \in L_0^2(S, \mathcal{S}, \mu)$ for $u \in S$, f_u being a constant vector for $u \notin S$, and $g_v \in L^2(E, \mathcal{E}, \nu)$ being unrestricted.

Given a linear operator L acting on $L^2(S, \mathcal{S}, \mu)$ and a vertex $u \in G_n$ we may consider “ L applied at u ”. More precisely we define an operator $L^{(u)}$ acting on $\mathcal{H}_n$ as being unitary equivalent, via the isomorphism (2.1), to the tensor product of L acting on the copy of $L^2(S, \mathcal{S}, \mu)$ corresponding to u together with the identity on all other factors. We consider the case that L is given by $P_{1^\perp}$ which is the orthogonal projection onto the subspace $L_0^2(S, \mathcal{S}, \mu)$. We define an operator on $\mathcal{H}_n$ via

$$(2.3) \quad A_n = \sum_{u \in G_n} P_{1^\perp}^{(u)}.$$

By considering its action on the generating vectors for each subspace $\mathcal{H}_S$ we find that the eigenvalues of A_n are $0, 1, 2, \dots, 2^n$, with the eigenvalue k having corresponding eigenspace $\bigoplus_{|S|=k} \mathcal{H}_S$. Here $|S|$ denotes the number of vertices belonging to S . On the other hand the operator $I - P_{\mathcal{K}_n}$, which commutes with A_n , has eigenvalues 0 and 1 with corresponding eigenspaces $\mathcal{K}_n = \mathcal{H}_\emptyset$ and $\bigoplus_{S \neq \emptyset} \mathcal{H}_S$. By decomposing $f \in \mathcal{H}_n$ according to the subspaces $\mathcal{H}_S$ we deduce the following inequality.

Proposition 3. *For every $f \in \mathcal{H}_n$,*

$$0 \leq \|(I - P_{\mathcal{K}_n})f\|^2 \leq (f, A_n f).$$

Proof of Theorem 1: Subcritical case. Because of the recursive structure endogeny is equivalent to $\mathcal{H}_0 \subset \mathcal{K}$. So fix $f \in \mathcal{H}_0$, which, in a slight abuse of notation, we also treat as an element of $L^2(S, \mathcal{S}, \mu)$. To prove f belongs to $\mathcal{K}$ it is enough, by virtue of Proposition 3, to prove that

$$(f, A_n f) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

We may express $(f, A_n f)$ using couplings of the tree-indexed process. Fix a vertex $u_n \in G_n$. Let Ω'_n be a copy of the finite configuration space Ω_n and consider the product space $\Omega_n \times \Omega'_n$ equipped with co-ordinate maps $(\epsilon_u, \epsilon'_u; u \in \Gamma_{n-1})$ and $(\xi_u, \xi'_u; u \in \Gamma_n)$. Let the probability measure $\tilde{m}_n$ on this product space have both marginals equal to m_n and be such that

$\tilde{m}_n$ is supported on the set where $\epsilon_u = \epsilon'_u$ for all $u \in \Gamma_{n-1}$ and $\xi_u = \xi'_u$ for all $u \in G_n$ except u_n ;

$(\epsilon_u; u \in \Gamma_{n-1})$, $(\xi_u; u \in \Gamma_n)$ and ξ'_{u_n} are independent under $\tilde{m}_n$.

It is easily verified that

$$(f, P_{1^\perp}^{(u_n)} f) = \frac{1}{2} \int_{\Omega_n \times \Omega'_n} (f \circ \xi_\emptyset - f \circ \xi'_\emptyset)^2 d\tilde{m}_n = (\mu \otimes \mu) P_n^{(2)} g,$$

where $P_n^{(2)}$ is the n -step transition matrix for $P^{(2)}$ and g is the function $g(x, x') = \frac{1}{2}(f(x) - f(x'))^2$ on S^2 . Summing over the possible choices of u_n we obtain

$$(f, A_n f) = 2^n (\mu \otimes \mu) P_n^{(2)} g.$$

Since g is zero on the diagonal of S^2 this quantity can also be expressed using $P^{(-)}$, and then, since the spectral radius of $P^{(-)}$ is less than $\frac{1}{2}$ by hypothesis, we obtain the desired convergence to zero as n tends to infinity. $\square$

3 Dynamics and non-endogeny

Let L be the operator on $L^2(S, \mathcal{S}, \mu)$ associated with a matrix $(L(x, x'); x, x' \in S)$,

$$Lf(x) = \sum_{x' \in S} L(x, x')f(x').$$

We use the transition probabilities $P^{(2)}$ to determine a new operator $\mathcal{P}L$ also acting on $L^2(S, \mathcal{S}, \mu)$, which is defined via its associated matrix via,

$$(3.1) \quad (\mu \otimes L)P^{(2)} = \mu \otimes (\mathcal{P}L).$$

Here $(\mu \otimes L)(x, x') = \mu(x)L(x, x')$ is treated as a row vector on S^2 and $P^{(2)}$ acts by matrix multiplication on the right. In a similar way we may define a quadratic superoperator $\mathcal{Q}$ by using the mapping $T^{(2)}$,

$$(3.2) \quad T^{(2)}(\mu \otimes L, \mu \otimes L) = \mu \otimes (\mathcal{Q}L).$$

Both superoperators $\mathcal{P}$ and $\mathcal{Q}$ arise when considering the isometric embedding of Hilbert spaces: $L^2(S, \mathcal{S}, \mu) \rightarrow L^2(S^2 \times E, \mathcal{S}^2 \times \mathcal{E}, \otimes^2 \mu \otimes \nu)$ given by $f \mapsto f \circ \phi$. Given L acting on $L^2(S, \mathcal{S}, \mu)$, it is easily verified that the new operator $\mathcal{P}L$ satisfies

$$(3.3) \quad (f \circ \phi, (L \otimes I \otimes I)g \circ \phi)_{L^2(S^2 \times E, \mathcal{S}^2 \times \mathcal{E}, \otimes^2 \mu \otimes \nu)} = (f, (\mathcal{P}L)g)_{L^2(S, \mathcal{S}, \mu)},$$

for all $f, g \in L^2(S, \mathcal{S}, \mu)$. Similarly L and $\mathcal{Q}L$ satisfy

$$(3.4) \quad (f \circ \phi, (L \otimes L \otimes I)g \circ \phi)_{L^2(S^2 \times E, \mathcal{S}^2 \times \mathcal{E}, \otimes^2 \mu \otimes \nu)} = (f, (\mathcal{Q}L)g)_{L^2(S, \mathcal{S}, \mu)},$$

for all $f, g \in L^2(S, \mathcal{S}, \mu)$. It is a consequence of the recursive structure that these two relations extend as is recorded in the following proposition whose proof we omit.

Proposition 4. *If $v \in G_{n+1}$ is a daughter of some $u \in G_n$ then, for all $f, g \in \mathcal{H}_n$,*

$$(f, L^{(v)}g) = (f, (\mathcal{P}L)^{(u)}g).$$

If $v_1, v_2 \in G_{n+1}$ are daughters of distinct $u_1, u_2 \in G_n$, then for $f, g \in \mathcal{H}_n$,

$$(f, L^{(v_1)}L^{(v_2)}g) = (f, (\mathcal{P}L)^{(u_1)}(\mathcal{P}L)^{(u_2)}g).$$

If $v_1, v_2 \in G_{n+1}$ are the two daughters of some $u \in G_n$, then for $f, g \in \mathcal{H}_n$,

$$(f, L^{(v_1)}L^{(v_2)}g) = (f, (\mathcal{Q}L)^{(u)}g).$$

Corresponding to the principal eigenvalue ρ of $P^{(-)}$ is a left eigenvector κ satisfying $\kappa P^{(-)} = \rho \kappa$. Of course κ is only determined up to a scalar multiple and we make some arbitrary choice. Considered as a function on $S \setminus S^\nearrow$, κ is symmetric since $P^{(-)}$ preserves the space of vectors with this symmetry. It is conceivable that $P^{(-)}$ is not irreducible in which case there may be some further freedom in choosing κ . This does matter so long as we always choose it, as we may, to be symmetric. Next we define a symmetric operator Q on $L^2(S, \mathcal{S}, \mu)$ from κ via

$$(3.5) \quad (f, Qg)_{L^2(S, \mathcal{S}, \mu)} = -\frac{1}{2} \sum_{(x, x') \in S^2 \setminus S^\nearrow} (f(x') - f(x))(g(x') - g(x))\kappa(x, x').$$

Q is the generator of an S -valued Markov process which jumps from x to x' at rate $Q(x, x') = \kappa(x, x')/\mu(x)$. We can always assume that $\mu(x) > 0$ for all $x \in S$ by deleting part of S if necessary.

Define an operator on $\mathcal{H}_n$ via

$$(3.6) \quad Q_n = (2\rho)^{-n} \sum_{u \in G_n} Q^{(u)}.$$

This operator is the generator of a Markov process taking values in the finite configuration space Ω_n . The coordinates ξ_u , with $u \in G_n$, evolve independently, each a copy of the process generated by Q but with their speed altered by the factor $(2\rho)^{-n}$. At any instant the ξ_u coordinates for $u \in \Gamma_{n-1}$ are determined from the ξ_u coordinates with $u \in G_n$ by application of the map ϕ with the innovations $(\epsilon_u; u \in \Gamma_{n-1})$ fixed for all time.

The significance of the family of generators Q_n is that they have a certain consistency property that manifests itself at the level of the corresponding forms. We introduce the forms $\mathcal{E}_n$ defined $\mathcal{H}_n$ via

$$(3.7) \quad \mathcal{E}_n(f, g) = -(f, Q_n g) \quad \text{for } f, g \in \mathcal{H}_n.$$

Lemma 5. *The operator Q satisfies $\mathcal{P}Q = \rho Q$, and consequently the forms $\mathcal{E}_n$ are consistent in the sense that for any $m \leq n$,*

$$\mathcal{E}_m(f, g) = \mathcal{E}_n(f, g) \text{ for all } f, g \in \mathcal{H}_m.$$

Proof. In view of the relation between $\mathcal{P}$ and $P^{(2)}$ given by (3.1), to prove the first assertion we must verify that

$$(\mu \otimes Q)P^{(2)} = \rho(\mu \otimes Q).$$

$(\mu \otimes Q)(x, x') = \kappa(x, x')$ for $x \neq x'$ and so the desired equality holds on $S \setminus S^\nearrow$ since there it becomes $\kappa P^{(-)} = \rho \kappa$. We deduce that the equality must also hold on the diagonal by observing that $\mathcal{P}$ preserves the class of operators satisfying $L1 = 0$.

Suppose that $f, g \in \mathcal{H}_m$ then using what we have just shown together with Proposition 4 we obtain

$$\begin{aligned} \mathcal{E}_{m+1}(f, g) &= -(f, Q_{m+1}g) = -(2\rho)^{-(m+1)} \sum_{v \in G_{m+1}} (f, Q^{(v)}g) \\ &= -2(2\rho)^{-(m+1)} \sum_{u \in G_m} (f, (\mathcal{P}Q)^{(u)}g) = -(2\rho)^{-m} \sum_{u \in G_m} (f, Q^{(u)}g) = \mathcal{E}_m(f, g), \end{aligned}$$

which proves the consistency of the forms. □

It follows from the consistency of the $\mathcal{E}_n$ just established that we can define a form $\mathcal{E}$ on the dense subspace $\bigcup_n \mathcal{H}_n$ of $\mathcal{H}$ via $\mathcal{E}(f, g) = \mathcal{E}_n(f, g)$ whenever $f, g \in \mathcal{H}_n$. However it is not necessarily true that $\mathcal{E}$ is closable.

Lemma 6. *If $2\rho > 1$ then for $f \in \mathcal{H}_m$ for some m ,*

$$\sup_{n \geq m} \|Q_n f\| < \infty.$$

Proof. Consider $f \in \mathcal{H}_n$. Expanding Q_{n+1} as a sum and using Proposition 4, plus $\mathcal{P}Q = \rho Q$, gives

$$\begin{aligned}
(f, Q_{n+1}^2 f) &= (2\rho)^{-2n-2} \sum_{u,v \in G_{n+1}} (f, Q^{(u)} Q^{(v)} f) \\
&= 2(2\rho)^{-2n-2} \sum_{u \in G_n} (f, (\mathcal{Q}Q + \mathcal{P}Q^2)^{(u)} f) + 4(2\rho)^{-2n-2} \sum_{\substack{u,v \in G_n \\ u \neq v}} (f, (\mathcal{P}Q)^{(u)} (\mathcal{P}Q)^{(v)} f) \\
&= 2(2\rho)^{-2n-2} \sum_{u \in G_n} (f, (\mathcal{Q}Q + \mathcal{P}Q^2)^{(u)} f) - (2\rho)^{-2n} \sum_{u \in G_n} (f, (Q^2)^{(u)} f) + (f, Q_n^2 f) \\
&= (2\rho)^{-2n} \sum_{u \in G_n} (f, \hat{Q}^{(u)} f) + (f, Q_n^2 f),
\end{aligned}$$

where the operator $\hat{Q}$ acting on $L^2(S, \mathcal{S}, \mu)$ is given by

$$\hat{Q} = 2(2\rho)^{-2} (\mathcal{Q}Q + \mathcal{P}Q^2) - Q^2.$$

But now we compare the operator $\sum_{u \in G_n} \hat{Q}^{(u)}$ with the number operator A_n defined by (2.3). Notice that the constant $1 \in L^2(S, \mathcal{S}, \mu)$ satisfies $\hat{Q}1 = 0$. Thus each subspace $\mathcal{H}_S$ of $\mathcal{H}_n$ is an invariant subspace for $\sum_{u \in G_n} \hat{Q}^{(u)}$ whose restriction to $\mathcal{H}_S$ has norm $|S| \|\hat{Q}\|$. Thus we have

$$\sum_{u \in G_n} (f, \hat{Q}^{(u)} f) \leq \|\hat{Q}\| (f, A_n f).$$

If we express $(f, A_n f)$ in terms of $P^{(-)}$ as we did in the previous section then we find that it is bounded by some constant times $(2\rho)^n$, and thus we deduce that, for an appropriate constant C ,

$$(f, Q_{n+1}^2 f) \leq (f, Q_n^2 f) + C(2\rho)^{-n}.$$

If $2\rho > 1$ then the desired conclusion follows. $\square$

Proof of Theorem 1: Supercritical case. The consistency of the forms $\mathcal{E}_n$ can be expressed in terms of the corresponding operators as $Q_m f = P_{\mathcal{H}_m} Q_n f$ for $f \in \mathcal{H}_m$ and $n \geq m$. Consequently if $\sup_{n \geq m} \|Q_m f\| < \infty$ then, as n tends to infinity, $Q_n f$ converges in $\mathcal{H}$ to some limit we denote by $\dot{Q}_\infty f$. We see that $-\dot{Q}_\infty$ is a positive symmetric operator with

$$\mathcal{E}(f, g) = - \left(f, \dot{Q}_\infty g \right) \quad \text{for } f, g \in \bigcup_n \mathcal{H}_n,$$

and consequently $\mathcal{E}$ is closable [see Theorem X.23 of [8]].

By construction the subspace $\mathcal{K}_n$ lies in the kernel of the operator Q_n for each n . Hence

$$\mathcal{E}(f, f) = 0 \text{ for all } f \in \bigcup_n \mathcal{K}_n.$$

Denoting the closure of $\mathcal{E}$ by $\bar{\mathcal{E}}$ we deduce that

$$\bar{\mathcal{E}}(f, f) = 0 \text{ for all } f \in \mathcal{K}.$$

But $\bar{\mathcal{E}}(f, f)$ cannot be identically zero on $\mathcal{H}$ since Q , and hence $\mathcal{E}_0$, was not zero, thus $\mathcal{K} \neq \mathcal{H}$. $\square$

4 The critical case

The critical case $2\rho = 1$ is endogenous, provided we impose two additional non-degeneracy conditions:

$$(4.1) \quad \mathcal{H}_0 \cap \mathcal{K}^\perp \text{ is trivial;}$$

$$(4.2) \quad P^{(-)} \text{ is irreducible.}$$

The first of these conditions can be thought of as analogous to the condition on a multitype branching process that every initial condition leads to a non-zero probability of eventual extinction. The example given in the next paragraph suggests that it is not possible to dispense with some condition of this type. The second condition is probably not essential, but without it the proof given below would be considerably more complicated.

The following example shows how it is possible for the process to be non-endogenous even if $2\rho = 1$. Let $S = \{-1, +1\}$ and $E = \{0, 1\}$ with $\nu(0) = \nu(1) = 1/2$ and $\mu(-1) = \mu(+1) = 1/2$. Suppose that

$$(4.3) \quad \phi(x_0, x_1, z) = \mathbf{1}(z = 0)x_0 + \mathbf{1}(z = 1)x_1.$$

$S^2 \setminus S^\nearrow = \{(-1, +1), (+1, -1)\}$ and the transition matrix $P^{(-)}$ is $1/2$ times the identity matrix, so plainly $2\rho = 1$. Furthermore $\xi_0 \in \mathcal{H}_0$ is orthogonal to every subspace $\mathcal{K}_n$ and hence to $\mathcal{K}$, in particular this shows that endogeny does not hold. However this example does not satisfy the strong symmetry condition we have assumed for ϕ , namely that $\phi(x_0, x_1, z) = \phi(x_1, x_0, z)$ for all $x_0, x_1 \in S$ and $z \in E$. I do not know whether there are any examples, with S finite, and this symmetry assumption upheld, but for which (4.1) fails.

The sequence of subspaces $\mathcal{H}_0 \cap \mathcal{K}_n^\perp$ is decreasing, and since $\mathcal{H}_0$ is finite-dimensional, (4.1) can only hold if there exists some m for which

$$(4.4) \quad \mathcal{H}_0 \cap \mathcal{K}_m^\perp \text{ is trivial.}$$

Now consider the quadratic form $(f, P_{\mathcal{K}_m} f)$. If the preceding condition holds then this quadratic form restricted to $f \in \mathcal{H}_0$ is positive definite, and using again the fact that $\mathcal{H}_0$ is finite-dimensional we deduce that there exists an $\epsilon > 0$ such that

$$(4.5) \quad (f, P_{\mathcal{K}_m} f) \geq \epsilon(f, f) \text{ for all } f \in \mathcal{H}_0.$$

This extends, see the final paragraph of this section, to

$$(4.6) \quad (f, P_{\mathcal{K}_{n+m}} f) \geq \epsilon^{|S|}(f, f) \text{ for all } f \in \mathcal{H}_S,$$

for a subset $S \subseteq G_n$, where $n \geq 1$ is arbitrary, whilst m is as above. Decomposing $f \in \mathcal{H}_n$ into its components in the subspaces $\mathcal{H}_S$ as S varies through subsets of G_n , and using (4.14) below, gives,

$$(4.7) \quad (f, P_{\mathcal{K}_{n+m}} f) \geq \sum_{S \subseteq G_n} \epsilon^{|S|}(f, P_{\mathcal{H}_S} f) = (f, P_{\mathcal{K}_n} f) + \sum_{\substack{S \subseteq G_n \\ S \neq \emptyset}} \epsilon^{|S|}(f, P_{\mathcal{H}_S} f)$$

Consequently for a fixed f ,

$$(4.8) \quad \sum_{\substack{S \subseteq G_n \\ \bar{S} \neq \emptyset}} \epsilon^{|S|} (f, P_{\mathcal{H}_S} f) \rightarrow 0,$$

as $n \rightarrow \infty$.

The criticality assumption that $2\rho = 1$ implies, in the presence of the additional condition (4.2), that the sequence of matrices $2^n P_n^{(-)}$ is bounded as n varies. Consequently, for a fixed f ,

$$(4.9) \quad (f, A_n f) = \sum_{S \subseteq G_n} |S| (f, P_{\mathcal{H}_S} f)$$

is also bounded. The only way that this is consistent with (4.8) is for

$$(4.10) \quad \sum_{\substack{S \subseteq G_n \\ \bar{S} \neq \emptyset}} (f, P_{\mathcal{H}_S} f) \rightarrow 0,$$

which proves endogeny.

A couple of the steps used above need amplification. Start by considering a generalization of (2.1) and (2.2). By decomposing the tree at level n we obtain a natural isomorphism

$$(4.11) \quad \mathcal{H}_{n+m} \cong \bigotimes^{G_n} L^2(\Omega_m, \mathcal{F}_m, m_m) \bigotimes^{\Gamma_{n-1}} L^2(E, \mathcal{E}, \nu).$$

Splitting $L^2(\Omega_m, \mathcal{F}_m, m_m)$ into the space of constants together its orthogonal complement $L_0^2(\Omega_m, \mathcal{F}_m, m_m)$, we obtain the decomposition

$$(4.12) \quad \mathcal{H}_{n+m} = \bigoplus_{S \subseteq G_n} \mathcal{H}_S(m),$$

where for each subset S the subspace $\mathcal{H}_S(m)$ is generated by vectors of the form $\otimes_{u \in G_n} f_u \otimes_{v \in \Gamma_{n-1}} g_v$ with $f_u \in L_0^2(\Omega_m, \mathcal{F}_m, m_m)$ for $u \in S$, f_u being a constant for $u \notin S$, and $g_v \in L^2(E, \mathcal{E}, \nu)$ being unrestricted. Notice that for each subset S of G_n the subspace $\mathcal{H}_S$ is included in $\mathcal{H}_S(m)$. There is a corresponding decomposition

$$(4.13) \quad \mathcal{K}_{n+m} = \bigoplus_{S \subseteq G_n} \mathcal{K}_S(m),$$

where $\mathcal{K}_S(m)$ is a subspace of $\mathcal{H}_S(m)$. The orthogonal projection $P_{\mathcal{K}_{n+m}}$ acts on $\mathcal{H}_{n+m}$ by projecting each subspace $\mathcal{H}_S(m)$ onto the corresponding subspace $\mathcal{K}_S(m)$. Accordingly if $f \in \mathcal{H}_{n+m}$ is decomposed as $f = \sum f_S$ with $f_S \in \mathcal{H}_S(m)$ then

$$(4.14) \quad (f, P_{\mathcal{K}_{n+m}} f) = \sum_S (f_S, P_{\mathcal{K}_{n+m}} f_S)$$

a fact that was used at (4.7).

Further examination reveals that $\mathcal{H}_S(m)$ is naturally isomorphic to a tensor product of $\mathcal{K}_n$ together with $|S|$ copies of $\mathcal{H}_m^0$, where the latter is the orthogonal complement of the space of constants in $\mathcal{H}_m$. Similarly $\mathcal{K}_S(m)$ is naturally isomorphic

to a tensor product of $\mathcal{K}_n$ together with $|S|$ copies of $\mathcal{K}_m^0$. The restriction of $P_{\mathcal{K}_{n+m}}$ to $\mathcal{H}_S(m)$ respects this tensor product structure, acting as the identity on the factor of $\mathcal{K}_n$ tensored with copies of the natural projection from $\mathcal{H}_m^0$ to $\mathcal{K}_m^0$ on the other factors. Similarly the restriction of $P_{\mathcal{H}_0}P_{\mathcal{K}_{n+m}}$ to $\mathcal{H}_S(m)$ is the tensor product of the identity on $\mathcal{K}_n$ with $|S|$ copies of $P_{\mathcal{H}_0}P_{\mathcal{K}_m}$ acting on $\mathcal{H}_m^0$. The inequality (4.6) follows from this and (4.5) since the smallest eigenvalue of a tensor product of operators is the product of the smallest eigenvalues.

5 Convergence to the dynamics

Throughout this section we work with the case $2\rho > 1$ and we make the additional assumption

$$(5.1) \quad P^{(-)} \text{ is a primitive matrix.}$$

According to Perron-Frobenius theory, under this condition, the limit of the rescaled n -step transition matrices $\rho^{-n}P_n^{(-)}$ exists and is given by

$$(5.2) \quad \lim_{n \rightarrow \infty} \rho^{-n}P_n^{(-)}((x, x'), (y, y')) = \theta(x, x')\kappa(y, y'),$$

where θ is the left eigenvector of $P^{(-)}$ corresponding to ρ , and, as before, κ is the right eigenvector. Here we normalize θ and κ so that

$$(5.3) \quad \sum_{(x, x') \in S^2 \setminus S^\nearrow} \theta(x, x')\kappa(x, x') = 1.$$

We may also normalize so that

$$(5.4) \quad \sum_{(x, x') \in S^2 \setminus S^\nearrow} \theta(x, x')\mu(x)\mu(x') = 1,$$

which fixes a choice of κ . We assume throughout this section that Q is defined by (3.5) with this choice of κ . We deduce that

$$(5.5) \quad \rho^{-n}(\mu \otimes \mu - \mu^\nearrow)P_n^{(2)} \rightarrow \kappa^\star,$$

where $\kappa^\star(x, x') = \kappa(x, x')$ when $x \neq x'$ and $\kappa^\star(x, x) = -\sum_{x' \neq x} \kappa(x, x')$. Using the relationship (3.1) between the superoperator $\mathcal{P}$ and $P^{(2)}$ this may be recast as

$$(5.6) \quad \lim_{n \rightarrow \infty} \rho^{-n}\mathcal{P}^n P_{1^\perp} = -Q.$$

Suppose that $f, g \in \mathcal{H}_m$ then a straightforward application of Proposition 4 gives

$$(5.7) \quad \frac{1}{(2\rho)^n}(f, A_n g) = \frac{1}{2^m \rho^n} \sum_{u \in G_m} (f, (\mathcal{P}^{n-m} P_{1^\perp})^{(u)} g).$$

Thus in view of (5.6) we deduce that

$$(5.8) \quad \frac{1}{(2\rho)^n}(f, A_n g) \rightarrow -(f, Q_m g) = \mathcal{E}(f, g), \quad \text{for all } f, g \in \mathcal{H}_m.$$

Recall that we proved that the form $\mathcal{E}$ is closable, and let Q_∞ be the self-adjoint operator associated with its closure. If $f \in \bigcup_m \mathcal{H}_m$ the limit $\check{Q}_\infty f = \lim_n Q_n f$ exists and defines a operator $\check{Q}_\infty$ with domain $\bigcup \mathcal{H}_m$. The self -adjoint operator Q_∞ is an extension (the Friedrichs extension) of $\check{Q}_\infty$.

Proposition 7. *If $f \in \mathcal{H}_m$ for some m , then as n tends to infinity*

$$(2\rho)^{-n} A_n f \rightarrow -Q_\infty f,$$

in the metric topology of $\mathcal{H}$.

Proof. Convergence in the metric topology is implied by weak convergence together with convergence of the norms. Consequently it is sufficient to verify that if $f \in \mathcal{H}_m$ for some m , then as n tends to infinity,

$$\|(2\rho)^{-n} A_n f\| \rightarrow \|Q_\infty f\|,$$

noting that weak convergence follows from this and (5.8).

We begin by computing $\|Q_\infty f\|$. We have, for $f \in \mathcal{H}_m$, and $n > m$,

$$\begin{aligned} (f, Q_n^2 f) &= (2\rho)^{-2n} \sum_{u \in G_n} (f, (Q^2)^{(u)} f) + (2\rho)^{-2n} \sum_{\substack{u, v \in G_n \\ u \neq v}} (f, Q^{(u)} Q^{(v)} f) \\ &= (2\rho)^{-2n} \sum_{u \in G_n} (f, (Q^2)^{(u)} f) + 2 \sum_{r=m}^{n-1} 2^{2(n-r-1)} (2\rho)^{-2n} \sum_{u \in G_r} (f, (\mathcal{Q} \mathcal{P}^{n-r-1} Q)^{(u)} f) \\ &\quad + 2^{2(n-m)} (2\rho)^{-2n} \sum_{\substack{u, v \in G_m \\ u \neq v}} (f, (\mathcal{P}^{n-m} Q)^{(u)} (\mathcal{P}^{n-m} Q)^{(v)} f) \\ &= (2\rho)^{-2n} \sum_{u \in G_n} (f, (Q^2)^{(u)} f) + 2 \sum_{r=m}^{n-1} (2\rho)^{-2(r+1)} \sum_{u \in G_r} (f, (\mathcal{Q} Q)^{(u)} f) \\ &\quad + (2\rho)^{-2m} \sum_{\substack{u, v \in G_m \\ u \neq v}} (f, Q^{(u)} Q^{(v)} f). \end{aligned}$$

Now as n tends to infinity $Q_n f$ converges to $Q_\infty f$ in the metric topology of $\mathcal{H}$, and hence the limit of the lefthandside above is $\|Q_\infty f\|^2$. Turning to the righthandside the first term tends to zero, and consequently we deduce that

$$\|Q_\infty f\|^2 = (2\rho)^{-2m} \sum_{\substack{u, v \in G_m \\ u \neq v}} (f, Q^{(u)} Q^{(v)} f) + 2 \sum_{r=m}^{\infty} (2\rho)^{-2r-2} \sum_{u \in G_r} (f, (\mathcal{Q} Q)^{(u)} f).$$

A similar calculation is valid for $\|A_n f\|^2$. If we denote $P_{1^\perp}$ by L and $\rho^{-r} \mathcal{P}^r P_{1^\perp}$ by L_r , then

$$\begin{aligned} (2\rho)^{-2n} \|A_n f\|^2 &= (2\rho)^{-2n} \sum_{u \in G_n} (f, L^{(u)} f) + 2 \sum_{r=m}^{n-1} (2\rho)^{-2r-2} \sum_{u \in G_r} (f, (\mathcal{Q} L_{n-r-1})^{(u)} f) \\ &\quad + (2\rho)^{-2m} \sum_{\substack{u, v \in G_m \\ u \neq v}} (f, L_{n-m}^{(u)} L_{n-m}^{(v)} f). \end{aligned}$$

Letting n tend to infinity we observe that since L_n tends to $-Q$ we obtain termwise convergence to the expression for $\|Q_\infty f\|^2$. To complete the proof we appeal to dominated convergence noting that since $\mathcal{Q} L_n 1 = 0$, we have the estimate

$$\sum_{u \in G_r} (f, (\mathcal{Q} L_{n-r-1})^{(u)} f) \leq K(f, A_r f),$$

where $K = \sup_n \|QL_n\| < \infty$. □

Lemma 8. For $f \in \mathcal{H}_n$,

$$\|Q_\infty f\| \leq \frac{K}{(2\rho)^n} \|A_n f\|,$$

where $K = \sup\{\|Q_\infty f\| : f \in \mathcal{H}_0 \text{ and } \|f\| = 1\}$.

Proof. Recall the natural isomorphism

$$\mathcal{H}_n \cong \bigotimes^{G_n} L^2(S, \mathcal{S}, \mu) \bigotimes^{\Gamma_{n-1}} L^2(E, \mathcal{E}, \nu).$$

Similarly

$$\mathcal{H} \cong \bigotimes^{G_n} L^2(\Omega, \mathcal{F}, m) \bigotimes^{\Gamma_{n-1}} L^2(E, \mathcal{E}, \nu).$$

The operator $Q_\infty : \mathcal{H}_n \rightarrow \mathcal{H}$ can be written as a sum

$$Q_\infty = \frac{1}{(2\rho)^n} \sum_{u \in G_n} Q_\infty^{(u)},$$

where $Q_\infty^{(u)}$ is unitary equivalent the tensor product of $Q_\infty : L^2(S, \mathcal{S}, \mu) \rightarrow L^2(\Omega, \mathcal{F}, m)$ on the factor corresponding to the node $u \in G_n$ and the identity on all other factors. It is easy to see from this structure, together with the fact that $Q_\infty 1 = 0$ that if $f \in \mathcal{H}_n$ decomposes as $f = \sum_{S \subseteq G_n} f_S$ with $f_S \in \mathcal{H}_S$ then

$$\begin{aligned} \|Q_\infty f\|^2 &= \sum_S \|Q_\infty f_S\|^2 = \frac{1}{(2\rho)^{2n}} \sum_S \sum_{u,v \in S} (Q_\infty^{(u)} f_S, Q_\infty^{(v)} f_S) \\ &\leq \frac{K^2}{(2\rho)^{2n}} \sum_S |S|^2 \|f_S\|^2 = \frac{K^2}{(2\rho)^{2n}} \|A_n f\|^2. \end{aligned}$$

□

Lemma 9. $\bigcup_m \mathcal{H}_m$ is a core for Q_∞ .

Proof. Recall that $\mathring{Q}_\infty$ is the restriction of Q_∞ to $\bigcup \mathcal{H}_m$. To verify the claimed result it suffices to show that, for some $\alpha > 0$, the range of $(\alpha - \mathring{Q}_\infty)$ is dense in $\mathcal{H}$. For this shows that α belongs to the resolvent set of the closure of $\mathring{Q}_\infty$, and then we apply the criterion of Theorem X.1 of [8].

Fix some $\alpha > 0$. Let $R_n^\alpha : \mathcal{H}_n \rightarrow \mathcal{H}_n$ be the α resolvent of Q_n . Given $f \in \bigcup_m \mathcal{H}_m$ let $v_n = R_n^\alpha f$ for n sufficiently large. Let $g \in \bigcup \mathcal{H}_m$, then for all sufficiently large n ,

$$(g, (\alpha - Q_\infty)v_n) = (g, (\alpha - Q_n)v_n) = (g, f).$$

Now suppose that we know that $(\alpha - Q_\infty)v_n$ is uniformly bounded in norm, then we deduce that, for any $g \in \mathcal{H}$,

$$(g, (\alpha - Q_\infty)v_n) \rightarrow (g, f) \text{ as } n \rightarrow \infty.$$

Thus the range of $(\alpha - \mathring{Q}_\infty)$ is weakly dense, and consequently norm dense in $\mathcal{H}$.

To verify the supposition that $(\alpha - Q_\infty)v_n$ is uniformly bounded in norm we note that since v_n are uniformly bounded in suffices to verify that $Q_\infty v_n$ are also. Using the previous lemma and the fact that R_n^α and A_n commute we have

$$\|Q_\infty v_n\| \leq \frac{K}{(2\rho)^n} \|A_n v_n\| = \frac{K}{(2\rho)^n} \|A_n R_n^\alpha f\| = \frac{K}{(2\rho)^n} \|R_n^\alpha A_n f\| \leq \frac{K}{\alpha(2\rho)^n} \|A_n f\|,$$

and the righthandside is bounded as n tends to infinity. $\square$

Proof of Theorem 2. We know that $\cup \mathcal{H}_n$ is a common core for Q_∞ and for $A_n P_{\mathcal{H}_n}$. Thus the convergence established at Proposition 7 implies strong resolvent convergence of $(2\rho)^{-n} A_n P_{\mathcal{H}_n}$ to $-Q_\infty$ (see Theorem VIII.25 of [7]). $\square$

Convergence of the generators in the strong resolvent sense implies that the semi-groups converge in the strong operator topology. From this fact we obtain the following corollary, which can also be expressed in terms of $T^{(2)}$.

Corollary 10. *For each fixed $t > 0$, as $n \rightarrow \infty$,*

$$\mathcal{Q}^n(P_1 + e^{-(2\rho)^{-n}t} P_{1^\perp}) \rightarrow M_t,$$

where M_t acting on $L^2(S)$ is defined by identifying $\mathcal{H}_0$ with $L^2(S)$ and then setting $(f, M_t g)_{L^2(S)} = (f, e^{tQ_\infty} g)$.

Another interpretation of this corollary is available in terms of the spectral measures $\mu_f^{(n)}$ defined at (1.10). The rescaled measures

$$(5.9) \quad \tilde{\mu}_f^{(n)}([0, x]) = \mu_f^{(n)}([0, (2\rho)^n x]) \quad \text{for } x \geq 0,$$

converge weakly towards a measure μ_f whose Laplace transform is given by

$$(5.10) \quad \int_0^\infty e^{-tx} \mu_f(dx) = (f, e^{tQ_\infty} f).$$

References

- [1] D. Aldous and A. Bandyopadhyay, *A survey of Max-type Recursive Distributional Equations*. Annals of Applied Probability, **15**:22, (2005), 1047-1110.
- [2] A. Bandyopadhyay, *Bivariate Uniqueness and Endogeny for Recursive Distributional Equations : Two Examples*, math.PR/0407175.
- [3] J. D. Biggins, *Lindley-type equations in the branching random walk*. Stochastic process. Appl., **75**, (1998), 105-133.
- [4] A. Caliebe and U. Rösler, *Fixed points with finite variance of a smoothing transformation*. Stochastic process. Appl., **107**, (2003), 105-129.
- [5] R. Durrett and T.M. Liggett, *Fixed points of the smoothing transformation*. Z. Wahrsch. verw. Gebiete, **64**, (1983), 275-301.

- [6] Q. Liu, *Fixed points of a generalized smoothing transform and applications to branching processes*. Adv. Appl. Probab. **30**, (1998),85-112.
- [7] M. Reed and B. Simon, *Methods of Mathematical Physics I: Functional Analysis*. Academic Press, (1980).
- [8] M. Reed and B. Simon, *Methods of Mathematical Physics II: Fourier Analysis and Self-Adjointness*. Academic Press, (1980).
- [9] L.C.G. Rogers and D. Williams, *Diffusions, Markov processes, and martingales. Volume 2: Itô Calculus*. Wiley, (1987).
- [10] B. Tsirelson, *Nonclassical stochastic flows and continuous products*. Probability Surveys, **1**, (2004), 173-298.
- [11] B. Tsirelson and A.V. Vershik, *Examples of nonlinear continuous tensor products of measure spaces and non-Fock factorizations*. Reviews in Mathematical Physics **10**:1 (1998), 81-145.